



Use of the Daftardar-Jafari polynomials in Optimal Homotopy Asymptotic Method for the solution of one Dimensional Heat and Advection-Diffusion Equations

S. SHAH, R. NAWAZ**, M. ULLAH*, J. ZADA**

Department of Mathematic, Xian Jiaotong University Xian, Shannxi, China

Received 24th December 2016 and Revised 1st February 2017

Abstract: In this paper Daftardar-Jafari (DJ) polynomials have been introduced in the Homotopy of Optimal Homotopy Asymptotic Method (OHAM) for finding the approximate solution of one dimensional heat and advection-diffusion equations. The results of proposed method are compared with exact solution to find absolute errors. The obtained results reveal that the proposed scheme is more reliable, explicit and effective.

Keywords: Optimal Homotopy Asymptotic Method with DJ-Polynomials (OHAM-DJ), Heat Equation, Advection-Diffusion Equation.

1. INTRODUCTION

Partial Differential Equations play a vital role in different fields of Engineering Sciences and Physics such as Thermodynamics, Heat Transform, fluid flow and Wave Propagation. Mostly Partial Differential Equations are non-linear and solution of such type of equations is difficult, so the researchers show their interest in finding the approximate solutions as the exact solutions are rare. Various techniques have been proposed to find the approximate solutions of nonlinear differential equations. The well-known methods for finding the approximate solution of differential equations are RBFM (Hon, and Mao 1999), ADM (Wazwaz, 2007), (Wazwaz, 2000), VIM (Noor and Mohyuddin, 2007 and 2008), HPM (Ganji, and Sadighi, 2006), (Liang, and Jeffrey, 2009), and HAM (Liao, 2004), OHAM was introduced by Marinca, *et al.*, 2008, for finding approximate solution of Differential Equations (Marinca, *et al.*, 2008), (Marinca, *et al.*, 2009), (Herisanu, *et al.*, 2008), (Haq *et al.*, 2010) (Ali, *et al.*, 2010) (Nawaz, *et al.*, 2010). This method combines the properties of both asymptotic methods and homotopy based methods. Recently, a new scheme of using Daftardar-Jafari polynomials in the homotopy of OHAM for the analytic solution of non-linear differential equation has been introduced by Ali, *et al.*, 2013, which gives us more accurate results than that of OHAM (Ali, *et al.*, 2013). The author has shown the efficiency of proposed scheme for Nonlinear Klein-Gordon and Advection equations (Shah, and Nawaz, 2016), (Shah, *et al.*, 2016), Here we use the Optimal Homotopy Asymptotic Method with Daftardar-Jafari (DJ) polynomials (OHAM-DJ) for finding approximate solution of following one dimensional advection-diffusion equation (Mohebbi and Dehghan, 2010).

$$\frac{\partial u}{\partial t} + \beta \frac{\partial u}{\partial s} = \alpha \frac{\partial^2 u}{\partial s^2}, \quad a \leq s \leq b; \quad t \geq 0, \quad (1.1)$$

subject to the initial condition:

$$u(x, 0) = \phi(x); \quad s \in [a, b], \quad (1.2)$$

and the boundary conditions are:

$$\begin{aligned} u(a, t) &= y_0(t), \\ u(b, t) &= y_1(t), \quad t \in [0, T]. \end{aligned} \quad (1.3)$$

For $\beta = 0$, the advection-diffusion equation reduces to one-dimensional heat equation. The Advection-Diffusion Equation is of primary significance in many physical systems, especially those involving fluid flow (Dehghan, 2004). It arises regularly in transferring heat energy, vorticity and mass in engineering and chemistry (Noye, 1990). and water transport in soils (Caglar, *et al.*, 2008)

2. FUNDAMENTAL MATHEMATICAL THEORY OF OHAM-DJ.

The basic mathematical theory of OHAM-DJ [18-20] is as follow. Let us consider the nonlinear differential equation

$$L(u(r)) = F(u(r)) + f(r), \quad r \in \Omega \quad B(u, \frac{\partial u}{\partial n}) = 0, \quad (2.1)$$

with boundary conditions

$$B(v(r, q), \frac{\partial v(r, q)}{\partial n}) = 0. \quad (2.2)$$

Her L is linear operator, F is Non-linear operator, B is Boundary operator is and f come with common meanings. In view of OHAM-DJ, we erect a homotopy as:

**Corresponding authors Murad. Ullah

*Department of Mathematic, Islamia Collage University Peshawar, KP, Pakistan

**Department of Mathematic, Abdul Wali Khan University Mardan, KP, Pakistan

$$(1-q)[L(v(r,q)) - f(r)] = h(q)[L(v(r,q)) - f(r) + F(v(r,q))]. \quad (2.3)$$

In Eq. (2.3) q is an embedding parameter such that $q \in [0,1]$, $h(q)$ is non-zero auxiliary function for $q \neq 0$, and it is 0 for $q = 0$. The unknown function $v(r,q)$ start from $v(r,0) = u_0(r)$ to $v(r,1) = u(r)$ as q starts from 0 to 1.

The auxiliary function $h(q)$ is chosen in the form,

$$h(q) = \sum_{i=1}^m q^i C_i. \quad (2.4)$$

The convergence control parameters $C_1, C_2, \dots$, are to be evaluated latter.

Next, we use Taylor's series to expand the function $v(r,q)$ about q ,

$$V(r,q) = u_0(r) + \sum_{m=1}^{\infty} u_m(r, C_1, C_2, \dots, C_m) q^m. \quad (2.5)$$

The nonlinear function $F(v(r,p))$ is decomposed as

$$F(v(r,p)) = F(u_0) + q[F(u_0 + u_1) - F(u_0)] + q^2[F(u_0 + u_1 + u_2) - F(u_0 + u_1)] + \dots$$

The expression on the right hand side

$$F(u_0), [F(u_0 + u_1) - F(u_0)], [F(u_0 + u_1 + u_2) - F(u_0 + u_1)], \dots$$

are DJ-Polynomials. Precisely, these polynomials are the terms of the Taylor's series of the nonlinear term. The convergence of these polynomials has been proved (Bhalekaret al., 2011).

For simplicity and convenience these polynomials are expressed as:

$$F_0 = F(u_0),$$

$$F_m = F\left(\sum_{i=0}^m u_i\right) - F\left(\sum_{i=0}^{m-1} u_i\right),$$

we can now express

$$F(v(r,q)) = F_0 + \sum_{i=0}^{m-1} q^i F_i. \quad (2.6)$$

Making use of Eq. (2.4), (2.5) and (2.6) into Eq. (2.3) and comparing the coefficient of same powers of θ , the following linear equations can be directly integrated:

0th order problem:

$$Lu_0 = f(r), \quad B(u_0, \frac{\partial u_0}{\partial x}) = 0. \quad (2.7)$$

1st order problem:

$$Lu_1 = (1 - C_1) Lu_0 - C_1(F_0 - f),$$

$$B(u_1, \frac{\partial u_1}{\partial x}) = 0. \quad (2.8)$$

2nd order problem:

$$Lu_2 = (1 - C_1) Lu_1 - C_1 F_1 + C_2(L - F_0 - f), \quad B(u_2, \frac{\partial u_2}{\partial x}) = 0, \quad (2.9)$$

although we can build higher order problems simply but solutions till second order problems are sufficient to generate tremendous results.

If the series converges in (2.5), for $q = 1$ then

$$\tilde{v}(r) = \tilde{u}(r) = u_0(r) + u_1(r, C_1) + u_2(r, C_1, C_2). \quad (2.10)$$

By substituting Eq. (2.10) into Eq.(2.1) the resulting residual is

$$R(r, C_1, C_2) = L(\tilde{u}(r)) - F(\tilde{u}(r)) - f(r), \quad (2.11)$$

If $R = 0$, then the exact solution is u . In other case, we reduce R over domain of the problem.

The Galerekin's method and the method of Least squares are followed here.

In method of Least Squares, first we build the function

$$J(C_1, C_2) = \int_{a_1}^{b_1} \int_{a_2}^{b_2} \dots \int_{a_n}^{b_n} R^2 dr, \quad (2.12)$$

and then, minimizing it, we have

$$\frac{\partial J}{\partial C_1} = \frac{\partial J}{\partial C_2} = 0, \quad (2.13)$$

and following the Galerekin's method, we explain the following system:

$$\int_{a_1}^{b_1} \int_{a_2}^{b_2} \dots \int_{a_n}^{b_n} R \frac{\partial u}{\partial C_1} dr = 0, \int_{a_1}^{b_1} \int_{a_2}^{b_2} \dots \int_{a_n}^{b_n} R \frac{\partial u}{\partial C_2} dr = 0. \quad (2.14)$$

3. APPLICATION OF OHAM-DJ

Example. 3.1: Solution of Heat Equation via OHAM-DJ:

Now we solve the heat equation is as follow

$$\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial s^2} = 0, \quad (3.1.1)$$

with the initial condition is

$$u_0(s, 0) = \sin(\pi s). \quad (3.1.2)$$

Exact solution is

$$w(s, t) = e^{-\pi^2 t} \sin(\pi s), \quad (3.1.3)$$

Applying the basic idea of OHAM-DJ mentioned in section 2, on Eq. (3.1.1) we have the following:

0th order problem:

$$\frac{\partial u_0(s, t)}{\partial t} = 0, \quad u_0(s, 0) = \sin(\pi s). \quad (3.1.4)$$

Its solution is

$$u_0(s, t) = \sin(\pi s). \quad (3.1.5)$$

1st order problem:

$$\frac{\partial u_1(s, t)}{\partial t} = \frac{\partial u_0(s, t)}{\partial t} + C_1 \frac{\partial u_0(s, t)}{\partial t} - C_1 \frac{\partial^2 u_0(s, t)}{\partial s^2}, \quad u_1(s, 0) = 0. \quad (3.1.6)$$

Its solution is

$$u_1(s, t) = C_1 \pi^2 t \sin(\pi s). \quad (3.1.7)$$

2nd order problem:

$$\frac{\partial u_2(s, t)}{\partial t} = C_2 \frac{\partial u_0(s, t)}{\partial t} + \frac{\partial u_1(s, t)}{\partial t} + C_1 \frac{\partial u_1(s, t)}{\partial t} - C_2 \frac{\partial^2 u_0(s, t)}{\partial s^2} - C_1 \frac{\partial^2 u_1(s, t)}{\partial s^2}, \quad u_2(s, 0) = 0. \quad (3.1.8)$$

Its solution is

$$u_2(s, t) = \frac{1}{2} \pi^2 (2C_1 t + 2C_1^2 t + 2C_2 t + C_1^2 \pi^2 t^2) \sin(\pi s). \quad (3.1.9)$$

3rd order problem:

$$\begin{aligned} \frac{\partial u_3(s, t)}{\partial t} &= C_2 \frac{\partial u_0(s, t)}{\partial t} + \frac{\partial u_1(s, t)}{\partial t} + C_1 \frac{\partial u_1(s, t)}{\partial t} - \frac{\partial u_2(s, t)}{\partial t} - C_2 \frac{\partial^2 u_0(s, t)}{\partial s^2} - C_1 \frac{\partial^2 u_1(s, t)}{\partial s^2}, \\ u_3(s, 0) &= 0. \end{aligned} \quad (3.1.10)$$

Its solution is

$$\begin{aligned} u_3(s, t) &= \frac{1}{6} \pi^2 (6C_1 t + 12C_1^2 t + 6C_1^3 t + 6C_2 t + 12C_1 C_2 t + 6C_3 t + 6C_1^2 \pi^2 t^2 + 6C_1^3 \pi^2 t^2 \\ &\quad + 6C_1 C_2 \pi^2 t^2 + C_1^3 \pi^4 t^3) \sin(\pi s). \end{aligned} \quad (3.1.11)$$

Adding equations (3.3.5), (3.3.7), (3.3.9) and (3.3.11),

$$u(s, t) = u_0(s, t) + u_1(s, t) + u_2(s, t) + u_3(s, t). \quad (3.1.12)$$

With the help of least square method, we obtained the values of unknown constants,

$$C_1 = -0.9983569043416364, \quad C_2 = 0.000001294430269822666,$$

$$C_3 = 4.065725267344232 \times 10^{(-9)}$$


and substituting the values of C_1, C_2, C_3 in Eq. (3.1.12), the approximate solution of heat equation is as follow:

$$\begin{aligned} u(s, t) &= \sin(\pi s) - 9.853387696948161t \sin(\pi s) + \frac{1}{2} \pi^2 (-0.0032782029295024273t \\ &\quad + 9.837197638403133t^2) \sin(\pi s) + \frac{1}{6} \pi^2 (-0.00002388862903046629t + \\ &\quad 0.09690421324098253t^2 - 96.92972218268893t^3) \sin(\pi s). \end{aligned} \quad (3.1.13)$$

The following table shows the Absolute errors of 3rd OHAM-DJ solution at different time levels. The absolute errors are very small which reveals the good agreement of OHAM-DJ solution with exact solution.

Table 3.1.1: Absolute errors of 3rd order OHAM-DJ solution for example 3.1 at different time level.

s, t	$t = 0.0001$	$t = 0.0005$	$t = 0.001$
0.1	2.51504×10^{-12}	2.11553×10^{-12}	2.38698×10^{-14}
0.2	4.78395×10^{-12}	4.024×10^{-12}	4.54081×10^{-14}
0.3	6.58462×10^{-12}	5.53857×10^{-12}	6.23945×10^{-14}
0.4	7.74059×10^{-12}	6.5109×10^{-12}	7.34968×10^{-14}
0.5	8.13893×10^{-12}	6.84597×10^{-12}	7.72715×10^{-14}
0.6	7.74059×10^{-12}	6.5109×10^{-12}	7.34968×10^{-14}
0.7	6.58462×10^{-12}	5.53857×10^{-12}	6.23945×10^{-14}
0.8	4.78395×10^{-12}	4.024×10^{-12}	4.54081×10^{-14}
0.9	2.51504×10^{-12}	2.11553×10^{-12}	2.38698×10^{-14}
1.0	9.9675×10^{-28}	8.38411×10^{-28}	9.44168×10^{-30}

Fig. 1: Comparison of OHAM-DJ solution (solid line) with exact solution (dashed line) for $t = 0.001$ **Example. 3.2.**

The advection-diffusion equation with $\beta = 1, \alpha = 0.1$ is as follow

$$\frac{\partial u}{\partial t} + \frac{\partial u}{\partial s} = 0.1 \frac{\partial^2 u}{\partial s^2}, \quad 0 < s < 1, \quad t > 0, \quad (3.2.1)$$

With initial condition

$$u(s, 0) = e^{5s} \left(\cos\left(\frac{\pi s}{2}\right) + 0.25 \sin\left(\frac{\pi s}{2}\right) \right). \quad (3.2.2)$$

Exact solution of Eq. (3.2.1) is

$$w(s, t) = e^{5(s-\frac{t}{2})} e^{\left(\frac{-\pi^2 t}{40}\right)} \left(\cos\left(\frac{\pi s}{2}\right) + 0.25 \sin\left(\frac{\pi s}{2}\right) \right). \quad (3.2.3)$$

Applying the proposed method OHAM-DJ as discussed in section 2, we obtain

0th order problem:

$$\frac{\partial u_0(s, t)}{\partial t} = 0, \quad u_0(s, 0) = e^{5s} \left(\cos\left(\frac{\pi s}{2}\right) + 0.25 \sin\left(\frac{\pi s}{2}\right) \right). \quad (3.2.4)$$

Its solution is

$$u_0(s, t) = 1.e^{5.s} \text{Cos}(1.5707963267948966s) + 0.25e^{5.s} \text{Sin}(1.5707963267948966s). \quad (3.2.5)$$

1storder problem:

$$\frac{\partial u_1(s, t)}{\partial t} = \frac{\partial u_0(s, t)}{\partial t} + C_1 \frac{\partial u_0(s, t)}{\partial t} + C_1 \frac{\partial u_0(s, t)}{\partial s} - 0.1C_1 \frac{\partial^2 u_0(s, t)}{\partial s^2}, \quad u_1(s, 0) = 0. \quad (3.2.6)$$

Its solution is

$$u_1(s, t) = t(2.7467401100272344C_1 e^{5.s} \text{Cos}(1.5707963267948966s) + 0.6866850275068086C_1 e^{5.s} \text{Sin}(1.5707963267948966s)). \quad (3.2.7)$$

2ndorder problem:

$$\begin{aligned} \frac{\partial u_2(s, t)}{\partial t} = & C_2 \frac{\partial u_0(s, t)}{\partial t} + \frac{\partial u_1(s, t)}{\partial t} + C_1 \frac{\partial u_1(s, t)}{\partial t} + C_2 \frac{\partial u_0(s, t)}{\partial s} - 0.1C_2 \frac{\partial^2 u_0(s, t)}{\partial s^2} \\ & - C_1 \left(\frac{\partial u_1(s, t)}{\partial s} + 0.1 \frac{\partial^2 u_0(s, t)}{\partial s^2} - 0.1 \left(\frac{\partial^2 u_0(s, t)}{\partial s^2} + \frac{\partial^2 u_1(s, t)}{\partial s^2} \right) \right), \quad u_2(s, 0) = 0. \end{aligned} \quad (3.2.8)$$

Its solution is

$$\begin{aligned} u_2(s, t) = e^{5s} (2.7467401100272344C_1 t \text{Cos}(1.5707963267948966s) + \\ 2.7467401100272344C_1^2 t \text{Cos}(1.5707963267948966s) + \\ 2.7467401100272344C_2 t \text{Cos}(1.5707963267948966s) + \\ 3.7722906160162113C_1^2 t^2 \text{Cos}(1.5707963267948966s) + \\ 0.6866850275068086C_1 t \text{Sin}(1.5707963267948966s) + \\ 0.6866850275068086C_1^2 t \text{Sin}(1.5707963267948966s) + \\ 0.6866850275068086C_2 t \text{Sin}(1.5707963267948966s) + \\ 0.9430726540040528C_1^2 t^2 \text{Sin}(1.5707963267948966s)). \end{aligned} \quad (3.2.9)$$

3rdorder problem:

$$\begin{aligned} \frac{\partial u_3(s, t)}{\partial t} = & C_3 \frac{\partial u_0(s, t)}{\partial t} + C_2 \frac{\partial u_1(s, t)}{\partial t} + \frac{\partial u_2(s, t)}{\partial t} + C_1 \frac{\partial u_2(s, t)}{\partial s} + C_3 \frac{\partial u_0(s, t)}{\partial s} \\ & - 0.1C_3 \frac{\partial u_0^2(s, t)}{\partial s^2} + C_2 \frac{\partial u_1(s, t)}{\partial s} - 0.1 \frac{\partial u_0^2(s, t)}{\partial s^2} + 0.1 \left(\frac{\partial u_0^2(s, t)}{\partial s^2} + \frac{\partial u_1^2(s, t)}{\partial s^2} \right) \\ & C_3 \left(\frac{\partial u_2(s, t)}{\partial s} - 0.1 \left(\frac{\partial u_0(s, t)}{\partial s} + \frac{\partial u_1(s, t)}{\partial s} + \frac{\partial u_2(s, t)}{\partial s} + \frac{\partial u_1^2(s, t)}{\partial s^2} \right) + \right. \\ & \left. 0.1 \left(\frac{\partial u_0^2(s, t)}{\partial s^2} + \frac{\partial u_1^2(s, t)}{\partial s^2} \right) \right), \end{aligned} \quad (3.2.10)$$

with boundary condition:

$$u_3(s, 0) = 0.$$

Whose solution is

$$u_3(s, t) = -e^{5s} t (-4.853429173528852C_1 \text{Cos}(1.5707963267948966s) -$$

$$\begin{aligned}
& 5.493480220054469C_1^2 \cos(1.5707963267948966s) - \\
& 2.746740110027235C_1^3 \cos(1.5707963267948966s) - \\
& 2.7467401100272344C_2 \cos(1.5707963267948966s) - \\
& 5.493480220054469C_1C_2 \cos(1.5707963267948966s) - \\
& 2.7467401100272344C_3 \cos(1.5707963267948966s) - \\
& 13.331108582108026C_1^2 t \cos(1.5707963267948966s) - \\
& 10.437844907070225C_1^3 t \cos(1.5707963267948966s) - \\
& 10.437844907070225C_1C_2 t \cos(1.5707963267948966s) - \\
& 6.102848442267399C_1^3 t^2 \cos(1.5707963267948966s) + \\
& 0.2887166941154069C_1 \sin(1.5707963267948966s) - \\
& 1.3733700550136172C_1^2 \sin(1.5707963267948966s) - \\
& 0.6866850275068087C_1^3 \sin(1.5707963267948966s) - \\
& 0.6866850275068086C_2 \sin(1.5707963267948966s) - \\
& 1.3733700550136172C_1C_2 \sin(1.5707963267948966s) - \\
& 0.6866850275068086C_3 \sin(1.5707963267948966s) + \\
& 0.7930297241612521C_1^2 t \sin(1.5707963267948966s) - \\
& 0.5465577919234269C_1^3 t \sin(1.5707963267948966s) - \\
& 0.5465577919234268C_1C_2 t \sin(1.5707963267948966s) + \\
& 0.36304109196625733C_1^3 t^2 \sin(1.5707963267948966s).
\end{aligned} \tag{3.2.11}$$

From equations (3.2.5), (3.2.7), (3.2.9), and (3.2.11), we obtain:

$$u(x, t) = u_0(s, t) + u_1(s, t) + u_2(s, t) + u_3(s, t) \tag{3.2.12}$$

The values of unknown constants are obtained by least square method, for $t = 0.0001$ the values of

$C_1 = -0.0007977074697195564$, $C_2 = -451.8410780151379$, and $C_3 = 901.9643760326104$.

Now putting the values of C_1, C_2 and C_3 in Eq. (3.2.12), we get the approximate solution of heat equation:

$$\begin{aligned}
u(s, t) = & e^{5s} \cos(1.5708s) + 0.25e^{5s} \sin(1.5708s) + t(-0.0021911e^{5s} \cos(1.5708s) - \\
& 0.000547774e^{5s} \sin(1.5708s)) - e^{5s} t(-1238.35 \cos(1.5708s) - 3.76219t \cos(1.5708s) \\
& + 3.09787 \times 10^{-9} t^2 \cos(1.5708s) - 309.588 \sin(1.5708s) - 0.196999t \sin(1.5708s) \\
& - 1.84284 \times 10^{-10} t^2 \sin(1.5708s)) + e^{5s} (-1241.09t \cos(1.5708s) + \\
& 2.40045 \times 10^{-6} t^2 \cos(1.5708s) - 310.273t \sin(1.5708s) + 6.00112 \times 10^{-7} t^2 \sin(1.5708s))
\end{aligned} \tag{3.2.13}$$

The following table shows the Absolute errors of 3rd OHAM-DJ solution at different time levels. The absolute errors reveal the good agreement of OHAM-DJ solution with exact solution.

Table 3.2: Absolute errors of 3rd order OHAM solution for example 3.2 at different time levels.

$x \cdot t$	$t = 0.0001$	$t = 0.0005$	$t = 0.001$
0.11	2.51504×10^{-12}	2.11553×10^{-12}	2.38698×10^{-14}
0.12	4.78395×10^{-12}	4.024×10^{-12}	4.54081×10^{-14}
0.13	6.58462×10^{-12}	5.53857×10^{-12}	6.23945×10^{-14}
0.14	7.74059×10^{-12}	6.5109×10^{-12}	7.34968×10^{-14}
0.15	8.13893×10^{-12}	6.84597×10^{-12}	7.72715×10^{-14}
0.16	7.74059×10^{-12}	6.5109×10^{-12}	7.34968×10^{-14}
0.17	6.58462×10^{-12}	5.53857×10^{-12}	6.23945×10^{-14}
0.18	4.78395×10^{-12}	4.024×10^{-12}	4.54081×10^{-14}
0.19	2.51504×10^{-12}	2.11553×10^{-12}	2.38698×10^{-14}
0.20	9.9675×10^{-28}	8.38411×10^{-28}	9.44168×10^{-30}

The (Fig. 3.2 (a)) shows the plot of Approximate solution and the (Fig. 3.2 (b)) shows the Exact solution, which are in close agreement.




Fig 3.2(a), 3D Approximate solution of $u(s, t)$




Fig 3.2 (b), 3D Exact solution of $w(s, t)$

4. CONCLUSION

In this paper, we applied the newly developed method namely OHAM-DJ to advection diffusion and heat equation. Table-3.1 gives absolute errors of OHAM-DJ solution for heat equation and Table3.2 gives absolute errors of OHAM-DJ solution for Advection Diffusion equation. All these results show good conformity with the exact solution. We concluded that among the existing numerical techniques, OHAM-DJ may be considered as a nice modification.

REFERENCES:

Bhalekar, S. and V. Daftardar-Gejji, (2011). "Convergence of the new iterative method". International Journal of Differential Equations.

Caglar, H., M. Ozer and N. Caglar, (2008). "The numerical solution of the one-dimensional heat equation by using third degree B-spline functions", Chaos, Solitons and Fractals. 38, 1197-1201.

Dehghan, M. (2004). "Weighted finite difference techniques for the one-dimensional advection-diffusion equation", Applied Mathematics and Computation. 147, 307-319.

Ganji, D. D., and A. Sadighi, (2006), "Application of He's homotopy perturbation method to nonlinear coupled systems of reaction diffusion equations", International Journal of Nonlinear Sciences and Numerical Simulation 7, 411-418.

Hon, Y. C., and X. Z. Mao, (1999). "A radial basis function method for solving options pricing models", Journal of Financial Engineering, 8, 31-50.

Herisanu, N., V. Marinca, T. Dordea and G. Madescu, (2008), "A new analytical approach to nonlinear vibration of an electric machine", Proc. Romanian. Acad. Series A: Mathematics, Physics, Technical Sciences, Information Science, 9, 229-236.

Haq, S., M. Idrees, and S. Islam, (2010). "Application of optimal Homotopy asymptotic method to eighth order initial and boundary value problems", International Journal of Applied Mathematics and Computation, 2(4), 73-80.

Javed A., S. Islam, S. Islam, G. Zaman, (2010) "The solution of multipoint boundary value problems by the Optimal Homotopy Asymptotic Method", Computers and Mathematics with Application, 59(6), 2000-2006

Javed A., S., I. A. Shah, S. Islam and H. Khan (2013) "Application of Optimal Homotopy Asymptotic Method with Daftardar-Jaffari Polynomials to Non-linear

- Differential Equations” World Applied Sciences Journal 28(10): 1456-1462.
- Liang, S., and D. J. Jeffrey, (2009), “Comparison of homotopy analysis method and homotopy perturbation method through an evolution equation”, Preprint submitted to Elsevier.
- Liao, S., (2004), “Introduction to the Homotopy Analysis Method”, Chapman and Hall/CRC, Florida.
- Noor M., A., and S. T., Mohyuddin, (2008), “Variational Iteration Method for Fifth-Order Boundary Value Problems Using He’s Polynomials”, Mathematical Problems in Engineering, doi: 10.1155/954794.
- Marinca, V., N. Herisanu, C. Bota and B. Marinca, (2009), “An Optimal Homotopy Asymptotic Method applied to steady flow of a fourth-grade fluid past a porous plate, Applied Mathematics Letters 22, 245-251
- Marinca V. and N. Herisanu, (2008), “Application of Optimal Homotopy Asymptotic Method for solving nonlinear equations arising in heat transfer, International s in Heat and Mass Transfer 35, 710-715.
- Marinca V. and N. Herisanu, (2009), “Determination of periodic solutions for the motion of a particle past a rotating parabola by means of the Optimal Homotopy Asymptotic Method “, Journal of Sound and Vibration doi: 10.1016/j.jsv.11.005.
- Marinca, V., N. Herisanu, and I. Nemes, (2008), “An Optimal Homotopy Asymptotic Method with application to thin film flow”, Central European Journal of Physics 6, 648-653.
- Mohebbi A. and M. Dehghan, (2010). “High-order compact solution of the one-dimensional heat and advection-diffusion equations”, Applied Mathematical Modelling. 34, 3071-3084.
- Noor M. A. and S. T. Mohyuddin, (2007), “Variational iteration technique for solving higher order boundary value problems”, Applied Mathematics and Computation, vol. 189, no. 2, 1929–1942.
- Nawaz, R., M. N. Khalid, S. Islam, and S. Yasin, (2010). “Solution of tenth order boundary value problems using optimal homotopy asymptotic method (OHAM)”, Canadian Journal on Computing in Mathematics, Natural Sciences, Engineering and Medicine, 1(2), 37-54.
- Noye, B. J. (1990). “Numerical solution of partial differential equations”, Lecture Notes.
- Shah, S., R. and N. Nawaz, (2016), “Optimal Homotopy Asymptotic Method with DJ-Polynomials to nonlinear Advection equations”. Appl. Environ. Biol. Sci. 6(5), 91-106.
- Shah, Z., and S. Shah, (2016) “Use of the Daftardar-Jafari Polynomial in Optimal Homotopy Asymptotic Method for the solution of linear and nonlinear Klein Gordon equations”. Appl. Environ. Biol. Sci. 6(4), 71-81.
- Wazwaz, A. M. (2007), “A comparison between the variational iteration method and adomian decomposition method”, Journal of Computational and Applied Mathematics, 207, 129–136.
- Wazwaz, A. M. (2000), “The Modified Adomian Decomposition Method for Solving Linear and Nonlinear Boundary Value Problems of Tenth-order and Twelfth-order”, Int. J. Nonlinear Sci. Num.Simul., 1, 17-24.