



Records of Carabidae in Fauna of British India Segregated for Present Boundaries of Pakistan

A. J. KAZI, I. KHATRI, M. A. RUSTAMANI, M.S.WAGAN*

Department of Entomology, Sindh Agriculture University Tadojam, Pakistan

Received 17th August 2015 and Revised 8th July 2016

Abstract: Present study is aimed to provide the comprehensive records of family carabidae of Pakistan, as mentioned in Fauna of British India. The study is further aimed to separate listed species, which were recorded from present boundaries of Pakistan. This is the first attempt to initiate and present an inventory of family carabidae, which is represented with 238 records within present boundaries of Pakistan.

Keywords: Ground Beetles, Fauna, Records, Pakistan

1. INTRODUCTION

Family carabidae consists of about 1500 genera and 40,000 species worldwide (Zhang, 2011), of which 238 carabid species are elucidated from Fauna of British India for present boundaries of Pakistan. Members of family carabidae are commonly called as ground beetles and are recognized with usually long, thread-like antennae, strongly margined pronotum and large prognathous mandibles. Carabids occupy most of the land habitats on nearly all continents. Bio-elements of Pakistan are largely Palaearctic (western Indus valley, Himalaya, Karakorum and more), secondly Oriental (Thar desert, southern Punjab, eastern Indus valley and more); lastly some elements are of Ethiopian (southwest of Balochistan) (Rafi *et al.*, 2010). Carabids are one of the most important insect groups used as an environmental indicators in many terrestrial ecosystems (Stork 1990, Desender *et al.* 1994). Despite their importance no comprehensive list of carabidae is available for Pak.

Indian Subcontinent remained under British rule during 1858 to 1947 (Burton, 2010), during ruling India they published a series of book famously known as *The fauna of British India, including Ceylon and Burma*. After the birth of Pakistan in 1947, the data of fauna found from present boundaries of Pakistan and reported within series

was needed to be separate. Looking at the importance of the family, the present attempt was undertaken and is presented here.

2. MATERIALS AND METHODS

For present studies five volumes of *the fauna of British India* were encountered.

Format adapted: All the species are presented in the genera mentioned in Fauna of British India. Each listed species is provided with page number and numbered reference with the help of software Endnote. The localities were further confirmed with help of geographical maps.

Distributions mentioned in original script are replaced with present names within Pakistan. In list only Pakistan is mentioned for those species where in original script no further distribution is given (except India) or are widely distributed.

3. RESULTS

Family carabidae consists of about 1500 genera and 40,000 species worldwide (Zhang, 2011), of which 204 carabid species are elucidated from Fauna of British India for present boundaries of Pakistan. Checklist of ground beetles (Carabidae) is presented here with 204 records elucidated from series *The fauna of British India, including Ceylon and Burma*.

*Department of Zoology, University of Sindh Jamshoro, Pakistan

S.#	Speceis	Locality
1.	<i>Acanthophorus modicus</i> (Gahan, 1894): 25[1]	Punjab: Lahore
2.	<i>Aeolesthes sarta</i> (Solsky, 1871): 129[1]	Baluchistan: Quetta
3.	<i>Aglaophis humerosus</i> (Chevrolat, 1863): 303[1]	Pakistan
4.	<i>Apatophysis kashmiriana</i> Semenov 1901: 70[1]	Punjab; Kashmir
5.	<i>Apatophysis modica</i> Gahan 1906: 71[1]	Baluchistan: Quetta
6.	<i>Asaphidion championi</i> Andrewes, 1924: 84[2]	Punjab
7.	<i>Asaphidion substriatum</i> Andrewes, 1925: 89[2]	Punjab
8.	<i>Axonya championi</i> Andrewes 1923: 41[2]	Punjab
9.	<i>Bembidion ajmonis</i> Netolitzky, 1935: 132[2]	Kashmir
10.	<i>Bembidion algidum</i> Andrewes, 1935: 187[2]	Kashmir
11.	<i>Bembidion aquilum</i> Andrewes, 1924: 126[2]	Punjab; Kashmir; Skardu
12.	<i>Bembidion babaulti</i> Andrewes, 1924: 176[2]	Punjab; Kashmir
13.	<i>Bembidion beesoni</i> Andrewes, 1933: 109[2]	Punjab
14.	<i>Bembidion betegara</i> Andrewes, 1924: 148[2]	Punjab
15.	<i>Bembidion bracculatum</i> Bates, 1889: 146[2]	Kashmir
16.	<i>Bembidion bucephalum</i> Netolitzky, 1920: 118[2]	Kashmir; Sindh
17.	<i>Bembidion cameroni</i> (Andrewes, 1922): 111[2]	Punjab
18.	<i>Bembidion caporiaccoi</i> Netolitzky, 1935: 204[2]	Kashmir; Sindh
19.	<i>Bembidion chitral</i> : 168[2]	Khyber Pakhtunkhwa: Chitral
20.	<i>Bembidion cimmerium</i> Andrewes, 1922: 128[2]	Punjab
21.	<i>Bembidion clops</i> Andrewes, 1935:174 [2]	Kashmir
22.	<i>Bembidion dardum</i> Bates, 1889: 163[2]	Kashmir
23.	<i>Bembidion discordans</i> Netolitzky, 1935b: 101[2]	Kashmir
24.	<i>Bembidion dux</i> Andrewes, 1935: 130[2]	Punjab
25.	<i>Bembidion eupages</i> Andrewes, 1934: 190[2]	Kashmir
26.	<i>Bembidion eutherum</i> Andrewes, 1923: 139[2]	Punjab
27.	<i>Bembidion exaratum</i> (Andrewes, 1924): 110[2]	Punjab
28.	<i>Bembidion exquisitum</i> Andrewes, 1923: 197[2]	Punjab
29.	<i>Bembidion fuscicrus</i> Motschulsky, 1855: 178[2]	Kashmir
30.	<i>Bembidion gagates</i> Andrewes, 1924: 142[2]	Pakistan
31.	<i>Bembidion gilgit</i> Fassati 1957: 171[2]	Gilgit; Kashmir
32.	<i>Bembidion hasurada</i> Fassati 1959: 138	Kashmir; Punjab
33.	<i>Bembidion hazara</i> Andrewes, 1935: 168[2]	Khyber Pakhtunkhwa: Hazara, upper Kagan Valley
34.	<i>Bembidion himalayanum</i> Andrewes, 1924: 119[2]	Khyber Pakhtunkhwa: Parachinar; Punjab
35.	<i>Bembidion holconotum</i> Andrewes, 1935: 138[2]	Kashmir
36.	<i>Bembidion hutchinsoni</i> Andrewes, 1934: 162[2]	Kashmir
37.	<i>Bembidion irregulare</i> Netolitzky 1934: 136[2]	Kashmir
38.	<i>Bembidion ixion</i> Andrewes, 1934: 166[2]	Kashmir
39.	<i>Bembidion kara</i> Andrewes, 1921: 114[2]	Punjab
40.	<i>Bembidion kempi</i> Andrewes, 1922: 150[2]	Punjab
41.	<i>Bembidion Khyber</i> Netolitzky, 1930: 162[2]	Khyber Pakhtunkhwa
42.	<i>Bembidion kurram</i> Andrewes, 1935: 120[2]	Khyber Pakhtunkhwa: Parachinar
43.	<i>Bembidion ladakense</i> Andrewes, 1924: 167 ^[2]	Punjab
44.	<i>Bembidion leander</i> Andrewes 1935: 127[2]	Punjab
45.	<i>Bembidion lissonotum</i> Bates, 1873: 192[2]	Kashmir
46.	<i>Bembidion loricatum</i> Andrewes, 1922: 103[2]	Punjab
47.	<i>Bembidion luntaka</i> Andrewes, 1924: 171[2]	Kashmir; Sindh; Punjab
48.	<i>Bembidion lysander</i> Andrewes, 1935: 144[2]	Punjab; Muree hills
49.	<i>Bembidion manicatum</i> Andrewes, 1935: 152[2]	Kashmir
50.	<i>Bembidion niloticum</i> Dejean, 1831: 199[2]	Punjab
51.	<i>Bembidion nivicola</i> Andrewes 1923: 121[2]	Kashmir
52.	<i>Bembidion orinum</i> Andrewes, 1922: 129[2]	Punjab
53.	<i>Bembidion pamiricola</i> Lutshnik 1928: 166[2]	Kashmir
54.	<i>Bembidion phaedrum</i> Andrewes, 1923: 146[2]	Punjab
55.	<i>Bembidion pluto</i> Andrewes 1924: 123[2]	Kashmir
56.	<i>Bembidion psilodorum</i> Andrewes, 1933: 180[2]	Kashmir
57.	<i>Bembidion radians</i> Andrewes 1922: 184[2]	Punjab
58.	<i>Bembidion regale</i> Andrewes, 1922: 129[2]	Punjab
59.	<i>Bembidion rubidum</i> Andrewes, 1924: 173[2]	Kashmir
60.	<i>Bembidion satanas</i> Andrewes, 1924: 127[2]	Khyber Pakhtunkhwa: Chitral; Sindh
61.	<i>Bembidion sillemi</i> Netolitzky, 1935: 175[2]	Sindh
62.	<i>Bembidion sobrinum</i> Dejean, 1831: 201[2]	Pakistan
63.	<i>Bembidion striaticeps</i> Andrewes, 1935: 104[2]	Punjab

64.	<i>Bembidion trimaculatum</i> : 202[2]	Punjab
65.	<i>Bembidion uvidum</i> Fassati 1955: 159 ^[2]	Punjab
66.	<i>Bembidion varium</i> Netolitzky 1929: 197 ^[2]	Kashmir
67.	<i>Bembidion xanthochiton</i> Andrewes, 1922: 154 ^[2]	Punjab
68.	<i>Bembidion xestum</i> Andrewes, 1923: 149 ^[2]	Punjab
69.	<i>Benibidion atalanta</i> : 178 ^[2]	Kashmir
70.	<i>Bernbidion leve</i> : 135 ^[2] Andrewes, 1924a: 78	Kashmir
71.	<i>Bernbidion limatum</i> : 135 ^[2]	Punjab
72.	<i>Brosicus punctatus</i> Dejean, 1828: 35 ^[2]	Peshawar
73.	<i>Caloclytus ignobilis</i> (Bates, 1878): 270 ^[1]	Murree
74.	<i>Calosoma beesoni</i> Andrewes 1919: 52[3]	Punjab
75.	<i>Calosoma himalayanicum</i> Gestro 1875: 55[3]	Kashmir; Ladakh Sindh; Punjab
76.	<i>Calosoma imbricatum</i> Brulle in d'Orbigny 1838: 63[3]	Sindh: Karachi
77.	<i>Calosoma maderae</i> , <i>F</i> , <i>var indicum</i> , Hope.: 62[3]	Khyber Pakhtunkhwa; Peshawar, Kohat; Punjab; Kashmir
78.	<i>Calosoma maderae</i> , <i>var kashmirensis</i> , Breun: 60[3]	Punjab; Kashmir
79.	<i>Calosoma maderae</i> Eidam 1926: 56[3]	Punjab
80.	<i>Calosoma maderae</i> , <i>var auropunctatum</i> , (Herbst, 1874): 56[3]	Kashmir; Sindh
81.	<i>Calosoma olivieri</i> Dejean 1831: 65[3]	Khyber Pakhtunkhwa; Peshawar, Baluchistan: Nushki and Quetta
82.	<i>Carabus barysomus</i> Sates, <i>var. htgeli</i> : 90[3]	Kashmir
83.	<i>Carabus barysomus</i> Bates, 1889: 87[3]	Kashmir, Khyber Pakhtunkhwa: Hazara, lower Kaghlan
84.	<i>Carabus barysomus</i> Bates, <i>var heroitcus</i> : 90[3]	Kashmir
85.	<i>Carabus boysi</i> Tatum, 1851: 95[3]	Kashmir; Punjab
86.	<i>Carabus caschmirensis</i> Redtenbacher in Huegel 1844: 91[3]	Kashmir; Punjab: Muree
87.	<i>Carabus dardiellus</i> Bates, 1889: 97[3]	Kashmir
88.	<i>Carabus dardiellus grannisparsus</i> : 99[3]	Kashmir; Punjab: Jhelum
89.	<i>Carabus dardiellus klagen</i> : 98[3]	Kashmir
90.	<i>Carabus dardiellus subpunctulus</i> Roeschke, 1907: 98[3]	Kashmir
91.	<i>Carabus epipleuralis</i> Semenov, 1906: 96[3]	Kashmir
92.	<i>Carabus rostianus</i> Semenov, 1906: 93[3]	Kashmir
93.	<i>Carabus stoliczkanus</i> Bates, 1878: 94[3]	Kashmir; Punjab: Muree
94.	<i>Ceresium rufum</i> Lameere, 1890: 157[1]	Pakistan
95.	<i>Chaetobrosicus anomalus</i> Chaudoir, 1878: 43[2]	Punjab; Kashmir; Gilgit
96.	<i>Cicindela albopunctata</i> Chaudoir 1852: 378[4]	Punjab
97.	<i>Cicindela aulica</i> Dejean, 1831: 374[4]	Sindh: Karachi
98.	<i>Cicindela bellana</i> Horn 1905: 433[4]	Sindh: Karachi
99.	<i>Cicindela chloris</i> (Hope, 1831): 376[4]	Kashmir; Gilgit
100.	<i>Cicindela chloropleura</i> (de Chaudoir, 1865): 327[4]	Punjab
101.	<i>Cicindela divesdives</i> Gory, 1833: 413[4]	Pakistan
102.	<i>Cicindela erudita</i> Wiedemann 1823: 362[4]	Kashmir; Punjab
103.	<i>Cicindela fastidiosa</i> (Dejean, 1825): 352[4]	Punjab: Rawal Pindi; Kashmir
104.	<i>Cicindela funebris</i> Schmidt-G. 1846: 341[4]	Punjab
105.	<i>Cicindela funerea</i> Horn 1926: 377[4]	Punjab
106.	<i>Cicindela germanica</i> (Linnaeus, 1758): 354[4]	Gilgit; Kashmir
107.	<i>Cicindela histrio</i> Tschitscherine 1903: 225[4]	Sindh: Karachi, Manora
108.	<i>Cicindela imperfecta</i> LeConte, 1851: 357[4]	Punjab
109.	<i>Cicindela intermedia</i> Chaudoir, 1852: 379[4]	Kashmir, Punjab
110.	<i>Cicindela leucoloma</i> : 352[4]	Punjab
111.	<i>Cicindela lunulata</i> Fabricius, 1781: 375[4]	Baluchistan: Noshki
112.	<i>Cicindela maindroni</i> (Horn): 432[4]	Sindh: Karachi, Island of Kiamari; Baluchistan
113.	<i>Cicindela mauritii</i> Horn, 1907: 392[4]	Pakistan
114.	<i>Cicindela melancholica</i> Fabricius, 1798: 356[4]	Sindh: Karachi
115.	<i>Cicindela nitida</i> Lichtenstein, 1796: 366[4]	Sindh: Karachi
116.	<i>Cicindela octonotata</i> Wiedemann 1819: 381[4]	Punjab
117.	<i>Cicindela ornata</i> Fleutiaux, 1898: 428[4]	Sindh: Karachi, on the sands of Manorah
118.	<i>Cicindela princeps</i> Vigers, 1825: 409[4]	Pakistan
119.	<i>Cicindela sublacerata</i> Solsky, 1874: 368[4]	Baluchistan; Kashmir; Skardu
120.	<i>Cicindela undulata</i> Latreille in Dejean 1825: 356[4]	Pakistan
121.	<i>Clivinaattenuata</i> (Herbst, 1806): 355[3]	Pakistan
122.	<i>Clivina gamma</i> Andrewes, 1929: 368[3]	Pakistan
123.	<i>Clivina lobata</i> Bonelli 1813: 375[3]	Pakistan
124.	<i>Clivina memnonia</i> Dejean 1831: 362[3]	Pakistan
125.	<i>Clivina striata</i> Putzeys 1846: 357[3]	Pakistan

126.	<i>Clivina tranquebarica</i> Bonelli 1813: 359[3]	Pakistan
127.	<i>Clytus monticola</i> Gahan 1906: 251[1]	Kashmir
128.	<i>Conosoma biguttatum</i> : 364[5]	India
129.	<i>Coryza semirubra</i> Andrewes 1926: 350[3]	Punjab
130.	<i>Craspedonotus schaum</i> Schaum, 1863: 39[2]	Khyber Pakhtunkhwa; Punjab
131.	<i>Cymbionotum candidum</i> Andrewes, 1935: 24[2]	Punjab
132.	<i>Cymbionotum luniferum</i> Andrewes 1935: 22[2]	Punjab: Faisalabad
133.	<i>Cymbionotum semeleleri</i> (Chaudoir, 1861): 21[2]	Baluchistan
134.	<i>Demonax ascendens</i> (Pascoe, 1859): 296[1]	Pakistan
135.	<i>Demonax lineatus</i> (Chevrolat, 1863): 281[1]	Pakistan
136.	<i>Derolus discicollis</i> Gahan 1906: 136[1]	Sindh: Karachi
137.	<i>Derolus mauritanicus</i> Buquet, 1840: 135[1]	Baluchistan
138.	<i>Diorthus simplex</i> Gahan, 1906: 133[1]	Baluchistan: Ormarah; Sindh: Karachi
139.	<i>Doesus telephoroides</i> Pascoe, 1862: 56[1]	Pakistan
140.	<i>Dyschirius hingstoni</i> Andrewes, 1929: 400[3]	Pakistan
141.	<i>Dyschirius indicus</i> Putzeys 1866: 406[3]	Pakistan
142.	<i>Dyschirius seistanus</i> Andrewes 1929: 394[3]	Pakistan
143.	<i>Hesperophanes cribricollis</i> Bates, 1878: 113[1]	Punjab: Murree
144.	<i>Hypoeshrus indicus</i> Gahan, 1906: 104[1]	Sindh: Karachi
145.	<i>Kunbir telephoroides</i> Lameere, 1890: 174[1]	Pakistan
146.	<i>Leistus kashmirensis</i> Andrewes, 1927: 110[3]	Kashmir
147.	<i>Leistus nivium</i> Andrewes 1925: 108[3]	Kashmir
148.	<i>Leptura rubriola</i> Bates, 1878: 81[1]	Kashmir; Punjab: Murree
149.	<i>Luperca laevigata</i> (Fabricius, 1781): 206[3]	Baluchistan
150.	<i>Macrotoma crenata</i> (Fabricius) Waterhouse, 1884: 36[1]	Kashmir
151.	<i>Megacephala euphratica</i> Dejean, 1822: 442[4]	Sindh: Karachi
152.	<i>Nebria barbata</i> Andrewes 1929: 130[3]	Kashmir
153.	<i>Nebria desgodinsi</i> Oberthur 1883: 127[3]	Punjab
154.	<i>Nebria himalayica</i> Bates 1889: 125[3]	Kashmir; Skardu; Punjab
155.	<i>Nebria limbiger a babaulti</i> Andrewes, 1924: 121[3]	Kashmir
156.	<i>Nebria limbiger a</i> Solsky, 1874: 120[3]	Kashmir; Punjab
157.	<i>Nebria masrina</i> Andrewes 1929: 128[3]	Kashmir
158.	<i>Nebria psammophila</i> Solsky, 1874: 118[3]	Kashmir; Punjab: Jhelam
159.	<i>Nebria xanthacra</i> Andrewes 1925: 122[3]	Punjab
160.	<i>Neocollyris attenuata</i> (Redtenbacher, 1848): 240[4]	Punjab
161.	<i>Neocollyris bipartita</i> (Fleutiaux, 1897): 264[4]	Pakistan
162.	<i>Neocollyris crassicornis</i> (Dejean, 1825): 261[4]	Pakistan
163.	<i>Neocollyris kollari</i> (Horn, 1901): 245[4]	Pakistan
164.	<i>Neocollyris redtenbacheri</i> (Horn, 1894): 239[4]	Punjab
165.	<i>Notiophilus orientalis</i> de Chaudoir 1850: 136[3]	Punjab: Muree; Kashmir; Khyber Pakhtunkhwa: Parachinar
166.	<i>Omophron axillaris</i> Chaudoir, 1868: 160[3]	Punjab
167.	<i>Omophron maculosus</i> : 151[3]	Pakistan
168.	<i>Omophron rotundatum</i> : 161[3]	Baluchistan: Quetta
169.	<i>Oxylobus alternans</i> Andrewes, 1929: 302[3]	Pakistan
170.	<i>Oxylobus meridionalis</i> Bates, 1891: 298[3]	Pakistan
171.	<i>Oxylobus porcatus</i> Andrewes 1929: 305[3]	Punjab
172.	<i>Oxylobus pygmaeus</i> Andrewes, 1929: 307[3]	South India
173.	<i>Oxymirus cursor</i> (Linnaeus, 1758): 79[1]	Baluchistan: Quetta
174.	<i>Paropisthius indicus</i> Chaudoir 1863: 104[3]	Pakistan
175.	<i>Paussus denticulatus</i> Westwood 1845: 494[4]	Pakistan
176.	<i>Paussus fulvus</i> Westwood 1841: 490[4]	Pakistan
177.	<i>Paussus saundersii</i> Westwood 1841: 483[4]	Pakistan
178.	<i>Paussus spencei</i> Westw.: 476[4]	Pakistan
179.	<i>Paussus stevensianus</i> Westwood, 1842: 489[4]	Pakistan
180.	<i>Perileptus ceylanicus</i> (Nietner, 1857): 51[2]	Punjab
181.	<i>Perileptus imaicus</i> Jeannel, 1923: 53[2]	Punjab
182.	<i>Platyrhopalus angustus</i> Westwood 1838: 461[4]	Pakistan
183.	<i>Pogonus micans</i> Chaudoir 1842: 310[2]	Baluchistan
184.	<i>Prionus corpulentus</i> Bates, 1878: 15[1]	Punjab: Murree; Kashmir
185.	<i>Prionus elliotti</i> , Gahan, 1906: 16[1]	Baluchistan: Quetta
186.	<i>Prothyma inornata</i> : 305[4]	Punjab
187.	<i>Prothyma limbata</i> : 304[4]	Punjab
188.	<i>Purpuricenus haussknechti</i> Witte, 1871: 186[1]	Kashmir; Punjab
189.	<i>Purpuricenus montanus</i> Plavil'shchikov 1940: 186[1]	Kashmir
190.	<i>Purpuricenus sanguinolentus</i> (Olivier, 1795): 184[1]	Pakistan

191.	<i>Remphan hopei</i> Waterhouse, 1836: 34[1]	Pakistan
192.	<i>Rhaphuma sexnotata</i> Chevrolat, 1863: 275[1]	Pakistan
193.	<i>Scapterus guerini</i> Dejean 1826: 330[3]	Punjab
194.	<i>Scarites cylindronotus</i> Faldermann, 1836: 238[3]	Baluchistan
195.	<i>Scarites dyschromus</i> Chaudoir, 1855: 243[3]	Pakistan
196.	<i>Scarites eurytus</i> Fischer v. W. 1828: 235[3]	Baluchistan: Noshki
197.	<i>Scarites guineensis</i> Dejean 1831: 274[3]	Sindh: Karachi
198.	<i>Scarites Indus</i> Olivier 1795[3]	Pakistan
199.	<i>Scarites liopterus</i> Chaudoir, 1880: 262[3]	Pakistan
200.	<i>Scarites mandibularis</i> Andrewes, 1929: 260[3]	Pakistan
201.	<i>Scarites nitens</i> Andrewes, 1929: 276[3]	Baluchistan
202.	<i>Scarites planus</i> Bonelli 1813: 279[3]	Baluchistan
203.	<i>Scarites punctum</i> Wiedemann 1823: 272[3]	Punjab
204.	<i>Scarites salinus</i> Dejean 1825: 239[3]	Baluchistan: Noshki
205.	<i>Scarites stenodes</i> Andrewes, 1929: 248[3]	Punjab: Multan
206.	<i>Scarites subcylindricus</i> de Chaudoir 1843[3]	Baluchistan
207.	<i>Scarites sulcatus</i> Olivier 1795: 242[3]	Punjab: Muree
208.	<i>Scarites tauropus</i> Andrewes, 1929: 244[3]	Pakistan
209.	<i>Scarites terricola</i> Bonelli, 1813 ^[3]	Baluchistan: Noshki
210.	<i>Siagona dilutipes</i> Andrewes 1929: 202[3]	Punjab
211.	<i>Siagona plicata</i> Andrewes, 1929: 185[3]	Pakistan
212.	<i>Siagona pubescens</i> Chaudoir, 1850: 186[3]	Pakistan
213.	<i>Syrdenus pallens</i> Andrewes, 1935 pallens: 311[2]	Sindh: Karachi
214.	<i>Tachys babaulti</i> (Andrewes, 1924): 271[2]	Punjab: Faisalabad
215.	<i>Tachys belli</i> : 251[2]	Sindh: Karachi
216.	<i>Tachys blandus</i> (Andrewes, 1924): 276[2]	Punjab
217.	<i>Tachys charis</i> Andrewes: 273[2]	Pakistan
218.	<i>Tachys comptus</i> Andrewes: 282[2]	Pakistan
219.	<i>Tachys haliploides</i> Andrewes 1925: 295[2]	Punjab
220.	<i>Tachys klugi</i> Nietner: 252[2]	Pakistan
221.	<i>Tachys latus</i> Peyr.: 249[2]	Kashmir; Sindh; Kaghan
222.	<i>Tachys ornatus</i> Burgeon 1935: 254[2]	Pakistan
223.	<i>Tachys poecilopterus</i> Bates, 1873: 259[2]	Punjab, Faisalabad
224.	<i>Tachys politus</i> Motschulsky: 268[2]	Pakistan
225.	<i>Tachys quadrillurn</i> : 228[2]	Punjab
226.	<i>Tachys stenseni</i> (Andrewes, 1923): 263[2]	Punjab
227.	<i>Tachys tetraspilus</i> (Solsky, 1874): 255[2]	Punjab
228.	<i>Tachys umbrosus</i> Motschulsky, 1851: 300[2]	Pakistan
229.	<i>Tachys vafer</i> : 264[2]	Pakistan
230.	<i>Tachys vilis</i> Andrewes, 1925: 221[2]	Kashmir
231.	<i>Tetropium oreinum</i> Gahan 1906: 95[1]	Kashmir
232.	<i>Trechus beasoni</i> Jeannel, 1930: 70[2]	Kashmir
233.	<i>Trechus cameroni</i> Jeannel 1923: 71[2]	Punjab
234.	<i>Trechus indicus</i> Jeannel 1928: 65[2]	Kashmir; Punjab: Muree
235.	<i>Trechus macrops</i> Jeannel, 1927: 67[2]	Kashmir
236.	<i>Trilophus interpunctatus</i> (Putzeys 1866): 383[3]	Punjab
237.	<i>Trinophyllin cribratum</i> : 156 ^[1]	Punjab: Muree; Kashmir
238.	<i>Xystrocera globosa</i> (Olivier, 1795): 106[1]	Pakistan

1. Andrewes, H., (1935) Carabidae 2. Harpalinae "The Fauna of British India, including Ceylon and Burma". 323-332.
2. Andrewes, H., (1995) Carabidae 1. Carabinae "The Fauna of British India, including Ceylon and Burma". 431-436
3. Fowler, W., (1912) Coleoptera, general introduction and cicindelidae paussidae "The Fauna of British India, including Ceylon and Burma".
4. **CONCLUSION**
 Checklist of ground beetles (Carabidae) is presented here with 204 records elucidated from series *The fauna of British India, including Ceylon and Burma*, all the records are presented in genera pre-
 sented by Fowler (1912) and Andrew (1929, 1935). Family of tiger beetles cicindelidae is included in present checklist and are treated here under Carabidae.

REFERENCES:

- Burton, S (2010). A History of India, John Wiley & Sons, London, 107-125,
- Cameron, M., (1932) Staphylinidae 3. (staphylinidae, trichophyinae, termitoridiscinae, pygosteninae, tachyporinae "The Fauna of British India, including Ceylon and Burma". 443-446.
- Desender, K., M. Dufrière and J. P. Maelfait (1994). Long term dynamics of carabid beetles in Belgium: a preliminary analysis on the influence of changing climate and land use by means of a database covering more than a century. In Carabid beetles: evolution 247-252. Springer Netherlands.
- Gahan, C., (2006) Coleoptera: cerambycidae "The Fauna of British India, including Ceylon 89-98.
- Kinnear N.B. (1991) "The history of Indian mammalogy and ornithology. Part I. Mammals." *Journal of Bombay Natural History Society* 50: 766-778.
- Rafi, M. A., W. Jürgen., M.A. Matin., A. Zia., A. Sultan and F. Naz. (2010). Faunistics of tiger beetles (Coleoptera: Cicindelidae) from Pakistan. *Journal of Insect Science*, 10(1), 116-122.
- Stork, N. E. (1990). The role of ground beetles in ecological and environmental studies. Intercept.
- Zhang, Z. Q. (2011). Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness: Magnolia press.